

2022

Bio prospecting of Lapachol producing endophytic fungi

Chandrashekar Srinivasa

Department of Studies in Biotechnology, Davangere University, Shivagangotri, Davangere, India

Sushma Pradeep

Department of Biotechnology and Bioinformatics, JSS Academy of Higher Education, Mysore, India

Shashank M Patil

Department of Biotechnology and Bioinformatics, JSS Academy of Higher Education, Mysore, India

Ramith Ramu

Department of Biotechnology and Bioinformatics, JSS Academy of Higher Education, Mysore, India

Shiva Prasad Kollur

Department of Sciences, Amrita School of Arts and Sciences, Amrita Vishwa Vidyapeetham, Mysuru Campus, Mysuru, India

See next page for additional authors

Follow this and additional works at: <https://rescon.jssuni.edu.in/ijhas>

 Part of the [Botany Commons](#)

Recommended Citation

Srinivasa, Chandrashekar; Pradeep, Sushma; Patil, Shashank M; Ramu, Ramith; Kollur, Shiva Prasad; Prasad, Shashanka K; and Shivamallu, Chandan (2022) "Bio prospecting of Lapachol producing endophytic fungi," *International Journal of Health and Allied Sciences*: Vol. 11: Iss. 1, Article 13.

DOI: 10.55691/2278-344X.1014

Available at: <https://rescon.jssuni.edu.in/ijhas/vol11/iss1/13>

This Review is brought to you for free and open access by Research Connect. It has been accepted for inclusion in International Journal of Health and Allied Sciences by an authorized editor of Research Connect.

Bio prospecting of Lapachol producing endophytic fungi

Cover Page Footnote

Acknowledgments: SC thank Davangere University, Davangere for the research support provided. SP, SMP, RR, SKP and CS acknowledge JSS AHER for the support and infrastructure provided. SPK acknowledges the Director Amrita Vishwa Vidyapeetham, Mysuru campus for laboratory infrastructural facilities.

Authors

Chandrashekar Srinivasa, Sushma Pradeep, Shashank M Patil, Ramith Ramu, Shiva Prasad Kollur, Shashanka K Prasad, and Chandan Shivamallu

REVIEW

Bioprospecting of Lapachol Producing Endophytic Fungi

Chandrashekar Srinivasa^a, M. Govindappa^{a,d}, Sushma Pradeep^b, Shashank M. Patil^b, Ramith Ramu^b, Shiva Prasad Kollur^c, Siddappa B. Kakkalameeli^{a,d}, Shashanka K. Prasad^b, Chandan Shivamallu^{b,*}

^a Department of Studies in Biotechnology, Davangere University, Shivangotri, Davangere, India

^b Department of Biotechnology and Bioinformatics, JSS Academy of Higher Education, Mysore, India

^c School of Physical Sciences, Amrita Vishwa Vidyapeetham, Mysuru Campus, Mysuru, India

^d Department of studies in Botany, Davangere University, Davangere, 577007, India

Abstract

Background: The association of endophytic fungi with medicinal plants has been one of the evolving areas of research in the past few decades. The secondary metabolites produced owing to such associations have been recognised for a wide range of biological activities.

Objectives: The aim of the present review is to highlight the isolation of lapachol from endophytic microorganisms with an emphasis on its biotransformation to improve its efficacy.

Methodology: The researchers followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) criteria. Published scientific articles on endophytes, host–pathogen interaction and lapachol isolation were collected from reputed journals from 1960 to 2022 using electronic databases using the key words as stated. Following that, the authors chose the required papers based on the criteria they devised. The data was extracted using the common research elements found in the publications.

Results and discussion: Lapachol is one such secondary metabolite known for potent antitumor properties. Synthesis of lapachol and its isolation from plant sources have been reported but an expensive process. Therefore, bioprospecting of this metabolite from endophytic fungi have been evaluated by a few researchers. It has been identified that *A. niger* and *Alternaria alternata* along with some of the filamentous endophytic fungi have been identified to produce lapachol. Some findings of biotransformation of lapachol to render it more potent have also been reported in the present review.

Conclusion: Lapachol is one of the plant secondary metabolites that possess several therapeutic benefits. Owing to its isolation being highly expensive from plant sources, endophytes have been exploited. Furthering to the isolation, the biotransformation of this bioactive molecule to enhance its efficiency has proven to be useful. In this regard, this review has enlightened some of the biotransformed lapachol and its derivatives with special emphasis on the endophytes that produce them. This review comprehensively highlights the various endophytic sources for lapachol production and its biotransformed derivatives.

Keywords: Endophytic, Secondary metabolites, Lapachol, Bioprospecting, *Alternaria alternata*

1. Introduction

Bioactive compounds have been increasingly popular in pharmaceuticals and naturopathy industry in recent years as a result of their human and plant health advantages. Although such

compounds from plant sources have been popular, microbial productions are gaining popularity for various benefits [1]. Microorganisms alone or in collaboration with their plant hosts are known to produce enumerable bioactives with noteworthy therapeutic abilities. Endophytes, or microbes that

Received 11 July 2022; revised 8 September 2022; accepted 14 September 2022.
Available online 16 December 2022

* Corresponding author.

E-mail addresses: chandru.s@davangereuniversity.ac.in (C. Srinivasa), sushmap@jssuni.edu.in (S. Pradeep), shashankmpatil@jssuni.edu.in, shashankaprasad@jssuni.edu.in (S.M. Patil), ramithramu@jssuni.edu.in (R. Ramu), shivachemist@gmail.com (S.P. Kollur), chandans@jssuni.edu.in (C. Shivamallu).

<https://doi.org/10.55691/2278-344X.1014>

2278-344X/© 2022 JSS Academy of Higher Education and Research. This is an open access article under the CC-BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

live inside plant tissues, create a variety of these chemicals as well [2]. Endophytes have synthesized a number of new antibiotics that are effective against multidrug-resistant bacteria and are under various levels of clinical trials. Several other therapeutic properties are also identified by such metabolites produced from endophytic origin. In addition, endophytic inoculation to plants promotes plant growth while also modulating the production of bioactive molecules with high medicinal potential [3]. This review will not exhaustively present the list of plant-associated microorganisms or the compounds produced by them, as these topics have already been covered. We also won't go over the precise methods for extracting and evaluating endophyte secondary metabolites that have already been discussed elsewhere [4]. Instead, this study will focus on some of the novel lapachol derivatives and their source endophytes, which are paving the way for new therapeutic development opportunities.

2. Endophytes – an insight

Endophytes are a class of microorganisms that inhabit plants naturally without adversely affecting the plant. Among the various endophytes, fungi are of great significance because they are a reservoir of indigenous bioactive compounds. They are ubiquitously present in several plant species capable of colonizing every part of the plant. The applications of endophytic fungi and their bioactives have drastically increased in the past decade with over 1000 research articles published only between 2010 and 2021. They are responsible for the assimilation of nutrients and defense against the host chemicals by synthesizing the required secondary metabolites [4]. In their struggle to survive amidst the host signaling, endophytes produce metabolites that are chemically similar to the ones found in the host plant owing to its symbiotic association [5,6]. Such interactions are more often controlled at the genetic level although certain abiotic factors also contribute to the production. Several studies have demonstrated the antibiotic [7], anti-inflammatory [8], antioxidant [9,10] and anticancer [11] properties of the compounds isolated from these endophytes.

Host plants have specific preferences to a distinctive endophyte that is suggestive of the divergence of endophytic composition. The diversity of fungal species within the host plant is based on the geographic location (temperate, tropic, aquatic and xerophytic) and specific plant part (leaves, twig, seeds and roots). Isolation of endophytic fungi from palm [12], Lichens [13], large trees [14,15], sea grasses [16] have been reported in

several studies. Furthermore, studies have been successfully performed to identify and characterize various bioactive metabolites from these endophytes that possess important biological activities. For instance, a breakthrough in this field of research was the identification of Taxol, an anticancer agent, from the endophyte *Taxomyces andreanae* [17]. Further studies on such isolation have led to the identification of an array of such secondary metabolites with bioactive properties. These products can be further scaled for large-scale production without hampering the biodiversity of medicinal plants [9]. An overview of various secondary metabolites and their biological activities are detailed in Table 1.

3. Host-endophyte interaction-a cost–benefit association

Plant-microbe compatibility is the key to a successful endophytic colonization. An endophyte is a microorganism that lives within a host plant, remains symptomless and unobtrusive without adversely affecting the plant [32]. Although the origin of endophytes is not clear, two theories entail the existence of an endophyte-an endogenous theory that states that endophytes were gaggled out of the mitochondria and chloroplasts of the plants and an exogenous theory that suggests that endophytes arrived from external environment and got associated with the host plant [33,34]. Yet, one of the most widely accepted phenomenon is the horizontal transmission of endophytic spores from a pool of colonizers that have the potential ability to symbiotically establish itself within the host [35]. Although most such associations are generalistic approach, some of the endophytes have specific affinity for a particular host. The host health is determined by the endophytic community within itself. For instance, stereotypical cacao-specific endophytic communities were established from leaf litter of healthy plants to its seedlings [36,37]. The endophytic community in this case was dominated by *C. tropicale* that was capable of suppressing pathogen damage to the host tissues. The specificity is dictated by the host secondary chemistry. However, in tropical plants, intra-species variation is often not remarkable and goes unnoticed [38]. Yet, the endophytic specificities prevail. The mechanism by which these chemicals are produced is not clearly understood. In addition, it is also not well understood whether these chemicals are produced by the host or the endophyte in response to its interaction with the host [39]. Some of the chemicals produced benefit the host by preventing pathogen attack while some others are produced to facilitate the host tissues

Table 1. Various secondary metabolites isolated from different endophytes with their corresponding biological activities.

Metabolite	Fungi	Host plant	Biological activity	Reference
Unspecified compound	<i>Colletotrichum gloeosporioides</i>	<i>Artemisia mongolica</i>	Antibiotic and fungistatic properties	[18]
Coronamycin	<i>Streptomyces</i> sp.	<i>Monstera</i> sp.	Antibiotic and antimalarial activity against <i>Plasmodium falciparum</i>	[19]
Camptothecin	<i>Fusarium solani</i>	<i>Camptotheca acuminata</i>	Anticancer	[20]
SZ-685C- an anthraquinone	<i>Halorosellinia</i> sp.	Mangrove	Antiproliferative against human breast, prostate, glioma and hepatoma cancer	[21]
Gliocladiocillins A and B	Unknown	Cordyceps-colonizing fungi	Apoptosis via intrinsic and extrinsic pathways in melanoma B16	[22]
anhydrofusarubin and methyl ether of fusarubin	<i>Cladosporium</i> sp.	<i>Cucumis sativus</i>	Cytotoxic against human leukemia cell line HL-60	[23]
Chaetoglobosin A	<i>Chaetomium globosum</i>	<i>Nymphaea nouchali</i>	Antimicrobial	[24]
Griseofulvin, cytochalasin D, multiploides, methyl coumarin, sordaricin, phomemone and xyloide	<i>Xylaria</i> sp.	Multiple hosts	Antifungal activity	[25,26]
Phomopsolide A	TC2-085 (unidentified endophyte)	<i>Geummacrophylum</i>	Anti-mycobacterial	[27]
Macrophorin A, ocomacrophorin and deacetoxyanthone A	<i>Gliomastix</i> sp.	<i>Phakelliafusca</i>	Anti-mycobacterial	[27]
Pullularins A	<i>Pullularia</i> sp.	<i>Quercus coccifera</i>	Antiviral	[28]
8-hydroxy-6,7-dimethoxy-3-methylisocoumarin	<i>Xylariaceae</i> sp.	<i>Quercus gilva</i> Blume	Alpha glucosidase inhibitory activity	[29]
7-methyl anthraquinone	<i>Penicillium</i> sp.	<i>Limonium tubiflorum</i>	Immunosuppressive	[30]
Lapachol	<i>Alternariaalternata</i>	<i>Tabebuia argentea</i>	Anticancer, anti-inflammatory, antiviral, antiparasitic	[31]

against biotic and abiotic stress. Therefore, studies have also suggested that the production of secondary metabolites is a defensive mechanism for the survival of the host plant and its endophyte [40].

One such natural bioactive compound produced from endophyte is lapachol (Fig. 1), known for its potential anticancer properties [9]. The first identification of this compound was from the red Lapacho tree ("Pau D'Arco"; *T. impetiginosa*). The traditional use of this tree for the treatment of several diseases such as cancer, malaria, mange trypanosomiasis, eczema, herpes, ulcers, syphilis, bacterial and fungal infections, skin disorders, fevers, stomach and bladder disorders led to several exploratory studies to assess the biological potential and chemical composition of this plant [40]. Lapachol was also isolated from various plants of the Bignoniaceae family including *Tabebuia flavescens*, *T. guayacan* [41], *Kigelia pinnata* [42], *Raderma cherasinica* [43], and *Melloya quadrivalvis* [44]. A synthetic analogue of lapachol obtained via a three-component system comprising of 2-hydroxy-1,4-naphthoquinone, 2-phenylindole and 4-nitrobenzaldehyde exhibited potent anticancer properties [45]. The study evaluated its activity against breast cancer and liver cancer both of which showed potent inhibitory activity against the growth of cancer cells with IC50 values of 6.52 and 12.97 $\mu\text{mol/L}$, respectively [46]. Furthermore, lapachol production via catalytic hydrogenation demonstrated significant anticancer properties against lung cancer, renal cancer, melanoma, breast cancer, prostate cancer and ovarian cancer [47]. Further studies carried out by Xue et al., demonstrated that lapachol was able to reduce the survival rate of rat brain microvascular endothelial cell [48]. Studies carried out by Cen et al., showed auxiliary properties of lapachol in interacting with the HIV-1 auxiliary protein Vpu thereby inhibiting the viral release [49].

4. Lapachol production from endophytic fungi

Isolation and biological assays on lapachol have been marked as one of the most interesting studies in phytochemistry. Dating back to 1952, Dr. Osvaldo




Fig. 1. Structure of the secondary metabolite-lapachol.

Goncalves de Lima evaluated the antitumor potential of lapachol based on its ability to produce xiloidone in presence of pyrimidine. Though synthetic production of lapachol and its isolation from plant sources have been well established, bioprospecting of this compound has recent come in light after the study carried out by Sadananda et al., [31]. Endophyte bioprospecting research is still in its infancy, with few formulations available at the pharmaceutical market [50]. This first ever report evaluated 13 endophytic fungi from various parts of *Tabebuia argentea* belonging to the bignoniaceae family. Among the different isolates, the study identified *Aspergillus niger* and *Alternaria alternata* as the two potent fungi capable of producing this naphthoquinone known as lapachol. This can be considered as a preliminary study for the isolation and characterization of lapachol-producing endophytes. Further, the study carried out by Govindappa, evaluated the antimitotic and antiproliferative potential of lapachol produced from endophytic fungal sources [51]. Among the various parts tested for isolation of fungal endophytes, leaf yielded *A. alternata*, *Alternaria* sp., *Rhizopus* sp., *Fusarium oxysporum*, *Fusarium* sp., *A. niger*, *Penicillium* sp., the bark yielded *A. alternata*, *F. oxysporum* and unidentified fungi whereas the stem yielded *A. alternata* and *Alternaria* sp. In this study, lapachol was confirmed to be produced by the leaf endophytes *A. alternata*, *Penicillium* sp., *Alternaria* sp., and *A. niger*, the bark endophytes were *A. alternata*. Furthermore, lapachol fraction from *A. niger* was proven to possess inhibitory potential against meristematic cells and therefore involved in inhibition of mitotic activity by a series of mechanisms such as chromosomal bridge and clumping, stickiness, abnormal distribution of metaphase chromosome, shrunken interphase cells, chromosomal degeneration, chromosomal bridges with vagrant chromosomes at anaphase, binucleolar and nucleolar burst and mega cell, large nucleolus and chromosomal clumping in small cells [52,53]. The extracted lapachol also demonstrated potent antiproliferative activity against yeast cells up to 70%. This study concluded that the anticancer properties of lapachol was remarkable and could be evaluated through further *in vitro* and *in vivo* experiments to uphold the druggability of the molecule.

In another interesting study *Mucorcircinelloides* NRRL3631, *Botrytis cinerea* UCA992 and *B. cinerea* 2100 were used for the biotransformation of lapachol. In this study, biotransformation into avicequinone-A by *M. circinelloides*, 3'-hydroxylapachol by *B. cinerea*, and into dehydro- α -lapachone by both *B. cinerea* and *M. circinelloides* were carried out. These

compounds were tested for their cytotoxic abilities that showed a higher cytotoxicity exerted by dehydro- α -lapachone over that of lapachol. This compound also revealed to be of lower toxicity to fungal cells suggesting the detoxification potential of the endophyte. A similar biotransformation study was carried out by Barbosa et al., that evaluated nine filamentous fungi namely, *A. flavus*, *A. fumigatus*, *F. oxysporum*, *Rhizopus microsporus*, *Penicillium cyclophilum*, *Paecilomyces variotii*, *Phanerochaete chrysosporium*, *Rhizomucor miehei*, and *Trichoderma asperellum* for their ability to transform lapachol into a less toxic and feasible metabolite [54]. The study opens avenues for using such filamentous fungi in the process of biotransformation to efficiently produce bioactive compounds with pharmaceutical significance.

5. Conclusion

The present review is on bioprospecting of lapachol producing endophytes. Lapachol is one of the well known secondary metabolite produced from plants belonging to the Bignoniaceae family. Further, an insight on the biotransformation of this secondary metabolite to render it less toxic and more potent has also been reported from a few studies that establish the biotransformed derivatives to be more potent than lapachol itself. In fact, our research shows that endophytic microbes constitute an underutilised bioresource with unique biological and chemical diversity. They have produced over 100 structurally distinct chemical compounds with a wide range of biological properties. In that, lapachol is one of the most potent metabolite with several biotransformed derivatives identified from endophytic origin. Yet, this review article indicates that only a few species have been studied thus far, implying that a large number of strains remain undiscovered. To fully understand the potential and diversity of bioactive metabolites, rigorous bioguided investigations are required. Investigating the metabolites produced by these and their subtle modifications to enhance their biological role could lead to the discovery of novel natural compounds with novel chemical frameworks for medicinal lead discovery.

Acknowledgments

SC thank Davangere University, Davangere for the research support provided. SP, SMP, RR, SKP and CS acknowledge JSS AHER for the support and infrastructure provided. SPK acknowledges the Director Amrita Vishwa Vidyapeetham, Mysuru campus for laboratory infrastructural facilities.

Conflict of interest

There is no conflict of interest.

References

- [1] Patil SM, Kumari VC, Sumana K, Sujay S, Tejaswini M, Shirahatti PS, et al. Sustainable development of plant tissue culture industry: the Indian scenario. *J Appl Biol Biotechnol* 2021 Mar;9(2):18–27. <https://doi.org/10.7324/JABB.2021.9202>.
- [2] Srinivasa C, Mellappa G, Patil SM, Ramu R, Shreevatsa B, Dharmashekar C, et al. Plants and endophytes—a partnership for the coumarin production through the microbial systems. *Mycology* 2022 Feb 6:1–4. <https://doi.org/10.1080/21501203.2022.2027537>.
- [3] Patil SM, Shirahatti PS, Ramu R. *Azadirachta indica* A. Juss (neem) against diabetes mellitus: a critical review on its phytochemistry, pharmacology, and toxicology. *J Pharm Pharmacol* 2021 Sep 25. <https://doi.org/10.1093/jpp/rgab098>.
- [4] Suto M, Takebayashi M, Saito K, Tanaka M, Yokota A, Tomita F. Endophytes as producers of xylanase. *J Biosci Bioeng* 2002 Jan 1;93(1):88–90. [https://doi.org/10.1016/S1389-1723\(02\)80059-7](https://doi.org/10.1016/S1389-1723(02)80059-7).
- [5] Newman DJ, Cragg GM. Natural products as sources of new drugs over the last 25 years. *J Nat Prod* 2007 Mar 23;70(3):461–77. <https://doi.org/10.1021/np068054v>.
- [6] Pandey PK, Singh S, Yadav RN, Singh AK, Singh MC. Fungal endophytes: promising tools for pharmaceutical science. *Int J Pharmaceut Sci Rev Res* 2014 Mar;25(2):128–38. <https://chfcau.org.in/sites/default/files/pdf/epub/FungalEndophytes-Pharmaceutical.pdf>.
- [7] Akinsanya MA, Ting A, Goh JK, Lim SP. Biodiversity, enzymatic and antimicrobial activities of bacterial endophytes in selected local medicinal plants. *J Biomed Pharmaceut Res* 2016;16(5):1.
- [8] Ting AS, Mah SW, Tee CS. Detection of potential volatile inhibitory compounds produced by endobacteria with biocontrol properties towards *Fusarium oxysporum* f. sp. *cubense* race 4. *World J Microbiol Biotechnol* 2011 Feb;27(2):229–35. <https://doi.org/10.1007/s11274-010-0447-y>.
- [9] Ejaz M, Javed S, Hamza M, Tabassum S, Abubakar M, Ullah I. Fungal endophytes are effective alternatives and novel sources of anticancer drugs. *Punjab Univ J Zool* 2020;35:13–24. <https://doi.org/10.17582/journal.pujz/2020.35.1.13.24>.
- [10] Mehanni MM, Safwat MS. Endophytes of medicinal plants. In: XIII International Conference on Medicinal and Aromatic Plants. 854; 2009 Oct 3. p. 31–9. <https://doi.org/10.17660/ActaHortic.2010.854.3>.
- [11] Hermanto A, Ting AS. Comparative effect of L-asparagine and sodium nitrate in inducing L-asparaginase production by endophytic *Fusarium* sp. *Acta Biol Szeged* 2016 Jan 1;60(2):145–50. <http://160.114.156.156/index.php/abs/article/view/2901>.
- [12] Patil MP, Patil RH, Maheshwari VL. A novel and sensitive agar plug assay for screening of asparaginase-producing endophytic fungi from aegle marmelos. *Acta Biol Szeged* 2012 Jan 1;56(2):175–7. <http://www.sci.u-szeged.hu/ABS>.
- [13] Suryanarayanan TS, Govindarajulu MB, Rajamani T, Tripathi M, Joshi Y. Endolichenic fungi in lichens of Champawat district, Uttarakhand, northern India. *Mycol Prog* 2017;16(3):205–11. <https://doi.org/10.1007/s11557-016-1268-7>.
- [14] Lamit LJ, Lau MK, Sthultz CM, Wooley SC, Whitham TG, Gehring CA. Tree genotype and genetically based growth traits structure twig endophyte communities. *Am J Bot* 2014 Mar;101(3):467–78. <https://doi.org/10.3732/ajb.1400034>.
- [15] Patil SM, Martiz RM, Ramu R, Shirahatti PS, Prakash A, Kumar BP, et al. Evaluation of flavonoids from banana pseudostem and flower (quercetin and catechin) as potent inhibitors of α -glucosidase: an in silico perspective. *J Biomol Struct Dyn* 2021 Aug 23:1–5. <https://doi.org/10.1080/07391102.2021.1971561>.
- [16] Murali TS. L-asparaginase from marine derived fungal endophytes of seaweeds. *Mycosphere* 2011;147–55. <http://eprints.manipal.edu/id/eprint/78455>.
- [17] Chow YY, Tan WS, Ting AS. Fungal endophytes from anticancer plants as producers of the antitumor agent L-asparaginase: a look at diversity, ubiquity, and enzyme production. In: Akhtar MS, Swamy MK, editors. *Anticancer plants: natural products and biotechnological implements*. Singapore: Springer; 2018. p. 233–53. https://doi.org/10.1007/978-981-10-8064-7_10.
- [18] Cosoveanu A, Rodriguez Sabina S, Cabrera R. Fungi as endophytes in *Artemisia thuscula*: juxtaposed elements of diversity and phylogeny. *J Fungi* 2018 Mar;4(1):17. <https://doi.org/10.3390/jof4020053>.
- [19] Ezra D, Castillo UF, Strobel GA, Hess WM, Porter H, Jensen JB, et al. Coronamycins, peptide antibiotics produced by a verticillate *Streptomyces* sp. (MSU-2110) endophytic on *Monstera* sp. *Microbiology* 2004 Apr 1;150(4):785–93. <https://www.microbiologyresearch.org/content/journal/micro/10.1099/mic.0.26645-0>.
- [20] Hriday M, Gorapi M, Hossain Z, Noor S, Chowdhury NS, Rahman M, et al. Putative anticancer compounds from plant-derived endophytic fungi: a review. *Molecules* 2022 Jan;27(1):296. <https://doi.org/10.3390/molecules27010296>.
- [21] Uzma F, Mohan CD, Hashem A, Konappa NM, Rangappa S, Kamath PV, et al. Endophytic fungi—alternative sources of cytotoxic compounds: a review. *Front Pharmacol* 2018 Apr 26;9:309. <https://doi.org/10.3389/fphar.2018.00309>.
- [22] Chen Y, Guo H, Du Z, Liu XZ, Che Y, Ye X. Ecology-based screen identifies new metabolites from a *Cordyceps*-colonizing fungus as cancer cell proliferation inhibitors and apoptosis inducers. *Cell Prolif* 2009 Dec;42(6):838–47. <https://doi.org/10.1111/j.1365-2184.2009.00636.x>.
- [23] Khan MI, Sohrab MH, Rony SR, Tareq FS, Hasan CM, Mazid MA. Cytotoxic and antibacterial naphthoquinones from an endophytic fungus, *Cladosporium* sp. *Toxicol Rep* 2016 Jan 1;3:861–5. <https://doi.org/10.1016/j.toxrep.2016.10.005>.
- [24] Dissanayake RK, Ratnaweera PB, Williams DE, Wijayarathne CD, Wijesundera RL, Andersen RJ, et al. Antimicrobial activities of endophytic fungi of the Sri Lankan aquatic plant *Nymphaea nouchali* and chaetoglobosin A and C, produced by the endophytic fungus *Chaetomium globosum*. *Mycology* 2016 Jan 2;7(1):1–8. <https://doi.org/10.1080/21501203.2015.1136708>.
- [25] Gehlot P, Singh J, editors. *Fungi and their role in sustainable development: current perspectives*. Singapore: Springer Singapore; 2018 Sep 9. <https://link.springer.com/book/10.1007/978-981-13-0393-7?noAccess=true>.
- [26] Mullin T, Clark T, Johnson J, Gray C. Isolation of phomopsolid A as the antimycobacterial constituent of an unidentified endophyte of the Canadian medicinal plant *Geum macrophyllum*. *Can J Urban Res* 2017 Jun 21;2(1). <https://ojs.library.ubc.ca/index.php/cjur/article/view/188630>.
- [27] He WJ, Zhou XJ, Qin XC, Mai YX, Lin XP, Liao SR, et al. Quinone/hydroquinone meroterpenoids with antitubercular and cytotoxic activities produced by the sponge-derived fungus *Gliomastix* sp. *ZSDS1-F7*. *Nat Prod Res* 2017 Mar 4;31(5):604–9. <https://doi.org/10.1080/14786419.2016.1207076>.
- [28] Isaka M, Berkaew P, Intereya K, Komwijit S, Sathitkunanon T. Antiplasmodial and antiviral cyclohexadepsipeptides from the endophytic fungus *Pullularia* sp. *BCC* 8613. *Tetrahedron* 2007 Jul 16;63(29):6855–60. <https://doi.org/10.1016/j.tet.2007.04.062>.
- [29] Indrianiingsih AW, Tachibana S. α -Glucosidase inhibitor produced by an endophytic fungus, *Xylariaceae* sp. *QGS 01* from *Quercus gilva* Blume. *Food Sci Hum Wellness* 2017 Jun 1;6(2):88–95. <https://doi.org/10.1016/j.fshw.2017.05.001>.
- [30] Lee JC, Lobkovsky E, Pliam NB, Strobel G, Clardy J. Subglutinols A and B: immunosuppressive compounds from the endophytic fungus *Fusarium subglutinans*. *J Org Chem* 1995 Nov;60(22):7076–7. <https://doi.org/10.1021/jo00127a001>.
- [31] Sadananda T, Nirupama S, Chaitra R, Govindappa KM, Chandrappa CP, Vinay Raghavendra B. Antimicrobial and

- antioxydant activities of endophytes from *Tabebuia argentea* and identification of Anticancer agent (Lapachol). *J Med Plants Res* 2011;5(16):3643–52. <https://doi.org/10.5897/JMPR.9000170>.
- [32] Whitaker BK, Reynolds HL, Clay K. Foliar fungal endophyte communities are structured by environment but not host ecotype in *Panicum virgatum* (switchgrass). *Ecology* 2018 Dec;99(12):2703–11. <https://doi.org/10.1002/ecy.2543>.
- [33] Fuentealba A, Pureswaran D, Bauce É, Despland E. How does synchrony with host plant affect the performance of an outbreaking insect defoliator? *Oecologia* 2017 Aug;184(4):847–57. <https://doi.org/10.1007/s00442-017-3914-4>.
- [34] Pushpa VH, Jayanthi MK, Patil SM, Ramu R. Pharmacological profile of Shiva Gutika: an uncharted and versatile polyherbal drug. *Life* 2021 Jan 1;14(1):215–9. <https://doi.org/10.1080/26895293.2021.1903570>.
- [35] Vincent JB, Weiblen GD, May G. Host associations and beta diversity of fungal endophyte communities in New Guinea rainforest trees. *Mol Ecol* 2016 Feb;25(3):825–41. <https://doi.org/10.1111/mec.13510>.
- [36] Patil SM, Shirahatti PS, VB CK, Ramu R, Prasad N. *Azadirachta indica* A. Juss (neem) as a contraceptive: an evidence-based review on its pharmacological efficiency. *Phytomedicine* 2021 Jul 15;88:153596. <https://doi.org/10.1016/j.phymed.2021.153596>.
- [37] Patil SM, Ramu R, Shirahatti PS, Shivamallu C, Amachawadi RG. A systematic review on ethnopharmacology, phytochemistry and pharmacological aspects of *Thymus vulgaris* Linn. *Heliyon* 2021 May 1;7(5):e07054. <https://doi.org/10.1016/j.heliyon.2021.e07054>.
- [38] Christian N, Herre EA, Clay K. Foliar endophytic fungi alter patterns of nitrogen uptake and distribution in *Theobroma cacao*. *New Phytol* 2019 May;222(3):1573–83. <https://doi.org/10.1111/nph.15693>.
- [39] Hartley SE, Eschen R, Horwood JM, Gange AC, Hill EM. Infection by a foliar endophyte elicits novel arabidopsid-based plant defence reactions in its host, *Cirsium arvense*. *New Phytol* 2015 Jan;205(2):816–27. <https://nph.onlinelibrary.wiley.com/doi/10.1111/nph.13067>.
- [40] Aghdam SA, Brown AM. Deep learning approaches for natural product discovery from plant endophytic microbiomes. *Environ Microb* 2021 Dec;16(1):1–20. <https://doi.org/10.1186/s40793-021-00375-0>.
- [41] Hamed AN, Mahmoud BK, Samy MN, Kamel MS. An extensive review on genus “*Tabebuia*”, family bignoniaceae: Phytochemistry and biological activities (1967 to 2018). *J Herb Med* 2020 Dec 1;24:100410. <https://doi.org/10.1016/j.hermed.2020.100410>.
- [42] Singh P, Khandelwal P, Sharma KK, Sharma MC. Cetyl triacantanoate and other constituents from *Acacia jacquemontii* and *Kigelia pinnata*. *J Indian Chem Soc* 2010 Nov 1;87(11):1403. https://www.researchgate.net/profile/Kuldeep-Sharma-11/publication/276271164_Cetyltriactanoate_and_other_constituents_from_Acacia_jacquemontii_Roxb_and_Kigelia_pinnata_DCJournal_of_Indian_Chemical_Society_Section_B_8711_1403-1407_2010/links/5554344308ae980ca60871c5/Cetyltriactanoate-and-other-constituents-from-Acacia-jacquemontii-Roxb-and-Kigelia-pinnata-DC-Journal-of-Indian-Chemical-Society-Section-B-8711-1403-1407-2010.pdf.
- [43] Inoue K, Chen CC, Inouye H, Kuriyama K. Quinones and related compounds in higher plants. Part 16. Naphthoquinones from *Radermachera sinica* Hemsl. (Bignoniaceae). *J Chem Soc* 1981;1:2764–70. <https://doi.org/10.1039/P19810002764>. *Perkin Transactions*.
- [44] Sanderman W, Simatupang MH, Wenderborn W. New quinones from the heartwood of *Paratecoma peroba*. *Naturwissenschaften* 1968;55:38.
- [45] Eyong KO, Krohn K, Hussain H, Folefoc GN, Nkengfack AE, Schulz B, et al. Newbouldiaquinone and newbouldiamide: a new naphthoquinone–anthraquinone coupled pigment and a new ceramide from *Newbouldia laevis*. *Chem Pharm Bull* 2005;53(6):616–9. <https://doi.org/10.1248/cpb.53.616>.
- [46] Zhang C, Qu Y, Jia Y, Long N, Bingxuan HF. α -Lapachol analog, its synthetic method and application as anticancer drug. 2016. <https://patents.google.com/patent/CN105503692A/en>.
- [47] Ottoni FM, Marques LB, Ribeiro JM, Franco LL, Souza Filho JD, Pádua RM, et al. Synthesis of lapachol-based glycosides and glycosyl triazoles with antiproliferative activity against several cancer cell lines. 2021. <https://doi.org/10.21203/rs.3.rs-916974/v1>.
- [48] Xue M, Guo C, Guan W. Application of lapachol in preparation of products for preventing and/or treating malignant glioma. 2013. CN103239430A, <https://patents.google.com/patent/CN103239430B/en>.
- [49] Cen S, Zhang Q, Mi Z. Application of lapachol for producing HIV-1 virus-resisting medicine. 2013. CN102885802A, <https://patents.google.com/patent/CN102885802A/en>.
- [50] Dos Santos IR, Abdel-Azeem AM, Mohesien MT, Piekutowska M, Sheir DH, da Silva LL, et al. Insights into the bioprospecting of the endophytic fungi of the medicinal plant *Palicourea rigida* Kunth (Rubiaceae): detailed biological activities. *J Fungi* 2021 Sep;7(9):689. <https://doi.org/10.3390/jof7090689>.
- [51] Govindappa M. First report of anti-cancer agent, lapachol producing endophyte, *Aspergillus Niger* of *Tabebuia argentea* and its in vitro cytotoxicity assays. *Bangladesh J Pharmacol* 2014 Mar 17;9(1):129–39. <http://bdpsjournal.org/index.php/bjp/article/view/180>.
- [52] Almeida ER. Preclinical and clinical studies of lapachol and beta-lapachone. *Open Nat Prod J* 2009 Apr 14;2(1). <https://doi.org/10.2174/1874848100902010042>.
- [53] Patil SM, Chandra SJ, Jayanthi MK, Shirahatti PS, Ramu R. Role of medicinal plants in the treatment of eumycetoma: a review. *J Appl Biol Biotechnol* 2021 Sep 1;9(5):1–8. <https://doi.org/10.7324/JABB.2021.9524>.
- [54] Barbosa CL, Fumagalli F, da Rosa-Garzon NG, da Silva Emery F, Cabral H. Lapachol biotransformation by filamentous fungi yields bioactive quinone derivatives and lapachol-stimulated secondary metabolites. *Prep Biochem Biotechnol* 2019 May 28;49(5):459–63. <https://doi.org/10.1080/10826068.2019.1591991>.